

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AGRICROP

AGRICROP

PROLAN 500TM

HERBICIDE

PROLAN 500TM

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L ORYZALIN

ACTIVE CONSTITUENT: 500 g/L ORYZALIN

GROUP D HERBICIDE

GROUP D HERBICIDE

For pre-emergent control of certain annual grasses and broadleaf weeds in non-bearing and bearing fruit and nut orchards, vineyards, nursery stock, ornamentals, amenity plantings, established and newly planted turf and industrial areas and rights of way as specified in the Directions for Use table.

For pre-emergent control of certain annual grasses and broadleaf weeds in non-bearing and bearing fruit and nut orchards, vineyards, nursery stock, ornamentals, amenity plantings, established and newly planted turf and industrial areas and rights of way as specified in the Directions for Use table.

IMPORTANT. READ THE ATTACHED BOOKLET BEFORE USE

IMPORTANT. READ THE ATTACHED BOOKLET BEFORE USE

AGRICROP PTY LTD ABN 84 100 473 309
255 Port Road, Hindmarsh SA 5007
Tel (07) 3346 4113 Fax (07) 3346 2702

AGRICROP PTY LTD ABN 84 100 473 309
255 Port Road, Hindmarsh SA 5007
Tel (07) 3346 4113 Fax (07) 3346 2702

APVMA Approval No. 60335/58307

APVMA Approval No. 60335/58307

NET CONTENTS

10L

NET CONTENTS

10L



AGRICROP



AGRICROP

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Agricrop Prolan 500 Herbicide

ACTIVE CONSTITUENT: 500 g/L ORYZALIN

GROUP	D	HERBICIDE
--------------	----------	------------------

For pre-emergent control of certain annual grasses and broadleaf weeds in non-bearing and bearing fruit and nut orchards, vineyards, nursery stock, ornamentals, amenity plantings and established and newly planted turf as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THIS CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING

APVMA Approval No: 60335/ 55477

Agricrop Pty Ltd
ABN: 84 100 473 309
255 Port Road
HINDMARSH SA 5007
Tel: 07 3348 4113
Fax: 07 3348 2702

**DIRECTIONS FOR USE IN ALL STATES
RESTRAINTS**

DO NOT use on newly seeded turf areas.

DO NOT use on turf under stress.

DO NOT use on Bent grass or Wintergrass.

DO NOT use on golf greens or bowling greens.

DO NOT apply if forecast heavy rain or storms are likely to cause runoff.

DO NOT irrigate to the point of runoff.

DO NOT apply by aircraft or through any type of irrigation equipment.

Crop Group				Weeds Controlled	Rate L/ha	Critical Comments
Banana ¹				Grasses: Barnyard Guinea grass Love grass Paradoxa grass Pigeon grass Spiny burr grass (Gentle Annie / Innocent Weed) Summer grass (Crabgrass ³) Broadleaf Weeds: Deadnettle Fathen Fumitory Portulaca (Pigweed) Sowthistle	Short term control (up to 4 months) 4.5L Long term control (6-8 months) 6.8L	See GENERAL INSTRUCTIONS ¹ DO NOT use as a pre-emergent at the time of planting with tissue culture banana planting material. ² Western Australia only: Vineyards on medium loam or heavier soil types, heavily infested with crabgrass may be sprayed with up to 9.9L/ha of Agricrop Prolan 500 and together with pome fruit and all Prunus species on these soils spraying can be immediately after transplanting from nurseries providing buds are still dormant and no functional leaves are present. ³ Western Australia only: Apply early in summer for crabgrass (<i>Digitaria sanguinalis</i>) control and incorporation with overhead irrigation.
Grape ²						
Pome Fruit	Apple	Pear	(see also non-bearing fruits)			
Stonefruit	Apricot	Peach	Nectarine			
	Cherry	Plum & prune				
Citrus	Grapefruit	Orange	(see also non-bearing fruits)			
Nuts	Almonds	Pecan	Walnut			
	Macadamia					
Non-bearing berryfruits	Blackberry	Currant	Raspberry			
	Blueberry	Gooseberry				
	Boysenberry	Loganberry				
Non-bearing – fruits including nursery stock of the following:	Avocado	Lemon	Olive			
	Custard apple	Litchi	Passionfruit			
	Feijoa	Mandarin	Pawpaw			
	Guava	Mango	Persimmon			
	Kiwifruit	Nashi	Rambutan			

Crop Group				Weeds Controlled	Rate L/ha	Critical Comments
Nursery stock and / or seedling stage conifers including species of the following genus:	<i>Abies</i> (fir trees) <i>Chamaecyparis</i> (Lawson and False cypress) <i>Cryptomeria cupressus</i> (cypress) <i>Juniperus</i> (Junipers)	<i>Picea</i> (Spruces) <i>Pinus</i> (pine) <i>Platycladus orientalis</i> (oriental arborvitae) <i>Podocarpus</i>	<i>Pseudotsuga menziessii</i> (Douglas fir or Oregon pine) <i>Sequoia sempervirens</i> (Coastal redwood) <i>Taxus cuspidata</i> (Japanese yew) <i>Thuja</i>	Wireweed (Hogweed) Brassica species: ⁴ Blackberry nightshade Caltrop Paddy melon Silverleaf nightshade		⁴ Control of these weeds may range from poor to excellent depending upon soil temperature, time of germination, depth of weed seed in the soil and amount and timing of soil moisture.

Crop Group		Weeds Controlled	Rate L/ha	Critical Comments
Nursery stock, ornamentals and amenity plantings comprising of the following:	<p>Australian native species of the genus:</p> <p><i>Acacia</i> (wattles) <i>Casuarina</i> (Oaks and sheoak) <i>Grevillea</i> <i>Agonis</i> (WA peppermint, Willow (or Honey) myrtle) <i>Chamaelucium uncinatum</i> (Geraldton wax) <i>Hakea</i> <i>Allocasuarina</i> <i>Clematis</i> <i>Hardenbergia</i> (sarsaparilla) <i>Banksia</i> <i>Correa</i> <i>Hibbertia</i> <i>Boronia</i> <i>Eriostemon myoporoides</i> (Wax flower) <i>Kennedia</i> <i>Callistemon</i> (bottlebrush) <i>Eucalyptus</i> <i>Lechenaultia</i> <i>Callitris</i> (Cypress pine) <i>Leptospermum</i> and <i>Melaleuca</i> (tea trees) <i>Calytrix</i> <i>Prostanthera</i> (mintbush) <i>Cassia</i> <i>Thryptomene</i> <i>Westringia</i></p>	<p>Grasses:</p> <p>Barnyard Guinea grass Love grass Pigeon grass Paradoxa grass Spiny burr grass (Gentle Annie / Innocent Weed) Summer grass (Crabgrass³)</p> <p>Broadleaf Weeds:</p> <p>Deadnettle Fathen Fumitory Portulaca (Pigweed) Sowthistle Wireweed (Hogweed)</p> <p>Brassica species:⁴</p> <p>Blackberry nightshade Caltrop Paddy melon Silverleaf nightshade</p>	<p>Short term control (up to 4 months) 4.5L</p> <p>Long term control (6-8 months) 6.8L</p>	<p>See GENERAL INSTRUCTIONS</p> <p>³ Western Australia only: Apply early in summer for crabgrass (<i>Digitaria sanguinalis</i>) control and incorporation with overhead irrigation.</p> <p>⁴ Control of these weeds may range from poor to excellent depending upon soil temperature, time of germination, depth of weed seed in the soil and amount and timing of soil moisture.</p>

Crop Group		Weeds Controlled	Rate L/ha	Critical Comments																																																																																							
Nursery stock, ornamentals and amenity plantings comprising of the following:	<p>Exotic species of the genus:</p> <table border="0"> <tr> <td><i>Abelia</i></td> <td><i>Fuschia</i></td> <td><i>Parahebe</i></td> </tr> <tr> <td><i>Acalypha</i></td> <td><i>Felicia</i></td> <td><i>Philadelphus</i></td> </tr> <tr> <td><i>Acer</i> (maples)</td> <td><i>Gardenia</i></td> <td><i>Philodendron</i></td> </tr> <tr> <td><i>Arctostaphylos</i></td> <td><i>Gingko</i></td> <td><i>Phoenix</i></td> </tr> <tr> <td><i>Ardisia</i></td> <td><i>Hibiscus</i></td> <td><i>Photinia</i></td> </tr> <tr> <td><i>Azalea</i></td> <td><i>Ilex</i></td> <td><i>Pieris</i></td> </tr> <tr> <td><i>Baccharis</i> (groundsel bush)</td> <td><i>Jasminum</i></td> <td><i>Pittosporum</i></td> </tr> <tr> <td><i>Bauhinia</i></td> <td><i>Justicia</i></td> <td><i>Populus</i> (poplar)</td> </tr> <tr> <td><i>Berberis</i> (barberry)</td> <td><i>Kalmia</i></td> <td><i>Prunus</i></td> </tr> <tr> <td><i>Betula</i> (birch tree)</td> <td><i>Koelreuteria</i></td> <td><i>Pseudopanax</i></td> </tr> <tr> <td><i>Bignonia</i></td> <td><i>Lagenaria</i></td> <td><i>Pyracantha</i></td> </tr> <tr> <td><i>Buxus</i> (box)</td> <td><i>Lagerstroemia</i></td> <td><i>Quercus</i></td> </tr> <tr> <td><i>Calluna</i></td> <td><i>Lantana</i></td> <td><i>Raphiolepis</i></td> </tr> <tr> <td><i>Camellia</i></td> <td><i>Lavendula</i> (lavender)</td> <td><i>Rhododendron</i></td> </tr> <tr> <td><i>Ceanothus</i></td> <td><i>Leucodendron</i></td> <td><i>Rosa</i> (roses)</td> </tr> <tr> <td><i>Cistus</i></td> <td><i>Leucothoe</i></td> <td><i>Rosemarinus</i> (rosemary)</td> </tr> <tr> <td><i>Clyostoma</i></td> <td><i>Ligustrum</i></td> <td><i>Russelia</i></td> </tr> <tr> <td><i>Coleonema</i></td> <td><i>Liquidambar</i></td> <td><i>Russiaea</i></td> </tr> <tr> <td><i>Coprosma</i></td> <td><i>Liriope</i></td> <td><i>Spiraea</i></td> </tr> <tr> <td><i>Cotoneaster</i></td> <td><i>Lonicera</i> (honeysuckle)</td> <td><i>Syringa</i></td> </tr> <tr> <td><i>Cuphea</i></td> <td><i>Magnolia</i></td> <td><i>Syzygium</i></td> </tr> <tr> <td><i>Cytisus</i> (broom)</td> <td><i>Mahonia</i></td> <td><i>Taxus</i></td> </tr> <tr> <td><i>Dodonea</i> (hopbush)</td> <td><i>Malus</i></td> <td><i>Trachelospermum</i> (<i>Rhymncospermum</i>)</td> </tr> <tr> <td><i>Erica</i> (heath)</td> <td><i>Metrosideros</i></td> <td><i>Umbellularia</i></td> </tr> <tr> <td><i>Escallonia</i></td> <td><i>Monstera</i></td> <td><i>Viburnum</i></td> </tr> <tr> <td><i>Euonymus</i></td> <td><i>Myrtus</i></td> <td><i>Washingtonia</i></td> </tr> <tr> <td><i>Eutaxia</i></td> <td><i>Nandina</i></td> <td><i>Weigela</i></td> </tr> <tr> <td><i>Fatsyhedera</i></td> <td><i>Nerium</i> (oleander)</td> <td><i>Xylosma</i></td> </tr> <tr> <td></td> <td><i>Osmanthus</i></td> <td><i>Yucca</i></td> </tr> </table>	<i>Abelia</i>	<i>Fuschia</i>	<i>Parahebe</i>	<i>Acalypha</i>	<i>Felicia</i>	<i>Philadelphus</i>	<i>Acer</i> (maples)	<i>Gardenia</i>	<i>Philodendron</i>	<i>Arctostaphylos</i>	<i>Gingko</i>	<i>Phoenix</i>	<i>Ardisia</i>	<i>Hibiscus</i>	<i>Photinia</i>	<i>Azalea</i>	<i>Ilex</i>	<i>Pieris</i>	<i>Baccharis</i> (groundsel bush)	<i>Jasminum</i>	<i>Pittosporum</i>	<i>Bauhinia</i>	<i>Justicia</i>	<i>Populus</i> (poplar)	<i>Berberis</i> (barberry)	<i>Kalmia</i>	<i>Prunus</i>	<i>Betula</i> (birch tree)	<i>Koelreuteria</i>	<i>Pseudopanax</i>	<i>Bignonia</i>	<i>Lagenaria</i>	<i>Pyracantha</i>	<i>Buxus</i> (box)	<i>Lagerstroemia</i>	<i>Quercus</i>	<i>Calluna</i>	<i>Lantana</i>	<i>Raphiolepis</i>	<i>Camellia</i>	<i>Lavendula</i> (lavender)	<i>Rhododendron</i>	<i>Ceanothus</i>	<i>Leucodendron</i>	<i>Rosa</i> (roses)	<i>Cistus</i>	<i>Leucothoe</i>	<i>Rosemarinus</i> (rosemary)	<i>Clyostoma</i>	<i>Ligustrum</i>	<i>Russelia</i>	<i>Coleonema</i>	<i>Liquidambar</i>	<i>Russiaea</i>	<i>Coprosma</i>	<i>Liriope</i>	<i>Spiraea</i>	<i>Cotoneaster</i>	<i>Lonicera</i> (honeysuckle)	<i>Syringa</i>	<i>Cuphea</i>	<i>Magnolia</i>	<i>Syzygium</i>	<i>Cytisus</i> (broom)	<i>Mahonia</i>	<i>Taxus</i>	<i>Dodonea</i> (hopbush)	<i>Malus</i>	<i>Trachelospermum</i> (<i>Rhymncospermum</i>)	<i>Erica</i> (heath)	<i>Metrosideros</i>	<i>Umbellularia</i>	<i>Escallonia</i>	<i>Monstera</i>	<i>Viburnum</i>	<i>Euonymus</i>	<i>Myrtus</i>	<i>Washingtonia</i>	<i>Eutaxia</i>	<i>Nandina</i>	<i>Weigela</i>	<i>Fatsyhedera</i>	<i>Nerium</i> (oleander)	<i>Xylosma</i>		<i>Osmanthus</i>	<i>Yucca</i>	<p>Grasses:</p> <p>Barnyard Guinea grass Love grass Paradoxa grass Spiny burr grass (Gentle Annie / Innocent Weed) Summer grass (Crabgrass³)</p> <p>Broadleaf Weeds:</p> <p>Deadnettle Fathen Fumitory Portulaca (Pigweed) Sowthistle Wireweed (Hogweed)</p> <p>Brassica species:⁴</p> <p>Blackberry nightshade Caltrop Paddy melon Silverleaf nightshade</p>	<p>Short term control (up to 4 months) 4.5L</p> <p>Long term control (6-8 months) 6.8L</p>	<p>See GENERAL INSTRUCTIONS</p> <p>³ Western Australia only: Apply early in summer for crabgrass (<i>Digitaria sanguinalis</i>) control and incorporation with overhead irrigation.</p> <p>⁴ Control of these weeds may range from poor to excellent depending upon soil temperature, time of germination, depth of weed seed in the soil and amount and timing of soil moisture.</p>
<i>Abelia</i>	<i>Fuschia</i>	<i>Parahebe</i>																																																																																									
<i>Acalypha</i>	<i>Felicia</i>	<i>Philadelphus</i>																																																																																									
<i>Acer</i> (maples)	<i>Gardenia</i>	<i>Philodendron</i>																																																																																									
<i>Arctostaphylos</i>	<i>Gingko</i>	<i>Phoenix</i>																																																																																									
<i>Ardisia</i>	<i>Hibiscus</i>	<i>Photinia</i>																																																																																									
<i>Azalea</i>	<i>Ilex</i>	<i>Pieris</i>																																																																																									
<i>Baccharis</i> (groundsel bush)	<i>Jasminum</i>	<i>Pittosporum</i>																																																																																									
<i>Bauhinia</i>	<i>Justicia</i>	<i>Populus</i> (poplar)																																																																																									
<i>Berberis</i> (barberry)	<i>Kalmia</i>	<i>Prunus</i>																																																																																									
<i>Betula</i> (birch tree)	<i>Koelreuteria</i>	<i>Pseudopanax</i>																																																																																									
<i>Bignonia</i>	<i>Lagenaria</i>	<i>Pyracantha</i>																																																																																									
<i>Buxus</i> (box)	<i>Lagerstroemia</i>	<i>Quercus</i>																																																																																									
<i>Calluna</i>	<i>Lantana</i>	<i>Raphiolepis</i>																																																																																									
<i>Camellia</i>	<i>Lavendula</i> (lavender)	<i>Rhododendron</i>																																																																																									
<i>Ceanothus</i>	<i>Leucodendron</i>	<i>Rosa</i> (roses)																																																																																									
<i>Cistus</i>	<i>Leucothoe</i>	<i>Rosemarinus</i> (rosemary)																																																																																									
<i>Clyostoma</i>	<i>Ligustrum</i>	<i>Russelia</i>																																																																																									
<i>Coleonema</i>	<i>Liquidambar</i>	<i>Russiaea</i>																																																																																									
<i>Coprosma</i>	<i>Liriope</i>	<i>Spiraea</i>																																																																																									
<i>Cotoneaster</i>	<i>Lonicera</i> (honeysuckle)	<i>Syringa</i>																																																																																									
<i>Cuphea</i>	<i>Magnolia</i>	<i>Syzygium</i>																																																																																									
<i>Cytisus</i> (broom)	<i>Mahonia</i>	<i>Taxus</i>																																																																																									
<i>Dodonea</i> (hopbush)	<i>Malus</i>	<i>Trachelospermum</i> (<i>Rhymncospermum</i>)																																																																																									
<i>Erica</i> (heath)	<i>Metrosideros</i>	<i>Umbellularia</i>																																																																																									
<i>Escallonia</i>	<i>Monstera</i>	<i>Viburnum</i>																																																																																									
<i>Euonymus</i>	<i>Myrtus</i>	<i>Washingtonia</i>																																																																																									
<i>Eutaxia</i>	<i>Nandina</i>	<i>Weigela</i>																																																																																									
<i>Fatsyhedera</i>	<i>Nerium</i> (oleander)	<i>Xylosma</i>																																																																																									
	<i>Osmanthus</i>	<i>Yucca</i>																																																																																									
Tea-Tree Oil plantations ⁵	Tea-Tree (<i>Melaleuca alternifolia</i>)		6.8L	⁵ Apply at time of planting and soon after harvesting. Apply in 200-450 L/ha water.																																																																																							

Crop Group	Weeds Controlled	Rate L/ha	Critical Comments
Turf Established Common couch Hybrid couch Kikuyu	Wintergrass (Poa annua) Annual ryegrass (Lolium spp)	4	Apply prior to germination of the weeds (usually from late summer to mid autumn, February to May). Apply every 16 weeks from early February to the end of August. The product should be incorporated by 10 – 15 mm of spray irrigation or rainfall as soon as possible, at least within one day of application. Allow at least 18 weeks between last application and overseeding/oversowing.
	Crabgrass Chickweed		Apply prior to germination of the weeds (usually from early spring, September to December) Apply every 16 weeks from early September to the end of February. The product should be incorporated by 10 – 15 mm of spray irrigation or rainfall as soon as possible, at least within one day of application. Allow at least 18 weeks between last application and overseeding/oversowing.

Crop Group	Weeds Controlled	Rate L/ha	Critical Comments
Turf Newly planted (i.e. line planted or sprigged) common couch, hybrid couch and kikuyu.	Wintergrass (<i>Poa annua</i>) Annual ryegrass (<i>Lolium</i> spp) Crabgrass Chickweed	2.4	Apply within the first 24 to 48 hours of the stolons being planted. The product should be incorporated by 10 – 15 mm of spray irrigation or rainfall as soon as possible, at least within one day of application. Apply every 8 weeks from planting; a maximum of 3 applications can be made from planting. Allow at least 10 weeks between last application and overseeding/oversowing. Note: the area that is being line planted or sprigged with stolons should be sprayed out with a non selective herbicide prior to planting.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

TURF: DO NOT graze treated turf/lawn or feed turf/grass clippings from any treated area to poultry or livestock.

GENERAL INSTRUCTIONS

Agricrop Prolan 500 Herbicide is a pre-emergent surface applied herbicide, which will control the specified annual grasses and broadleaf weeds in the situations listed.

Soil Preparation: Areas to be treated should be free of established weeds. Remove or thoroughly mix trash (weed residue, prunings etc.) into the soil before applying the product.

Soil Activation: Moisture in the form of rain or sprinkler irrigation (10-15mm) is required within 21 days of application to activate the product otherwise efficacy may be reduced. If moisture is not received within the period, incorporate the product into the top 2.5cm of soil(excluding established turf and newly planted turf).

Turf: Mechanical incorporation is not possible. Moisture in the form of rain or sprinkler irrigation (10-15mm) is therefore required within 21 days of application to activate the product otherwise efficacy may be reduced.

MIXING

The required amount of Agricrop Prolan 500 Herbicide should be added to half-full spray tank and agitated well during completion of filling. Do not leave spray solutions to stand over night.

APPLICATION

Apply Agricrop Prolan 500 Herbicide in 200 to 450 litres water per hectare. Apply uniformly, especially within the dripline of trees and shrubs, so that application above label rates does not occur. Use a properly calibrated low pressure (170 to 340 kPa) herbicide sprayer. Provide good by-pass or other agitation of the spray suspension in the tank before and during application. Soil treated with this product may be shallow cultivated without loss of herbicidal activity.

Turf: Ground Application: Spray nozzles should be uniformly spaced and of the same size, and should provide accurate and uniform application. Use spray nozzles that produce a **MEDIUM** spray quality. To ensure accuracy, calibrate sprayer before use and calibrate frequently. Apply at a volume of 200 to 450 L of water per hectare. Higher volumes should be used for severe weed infestations to ensure adequate spray coverage.

Good even coverage with the spray is essential for optimum weed control. Observe sprayer nozzles frequently during the spraying operation to ensure that the spray pattern is uniform. Avoid overlapping of spray runs. Avoid application under conditions when uniform coverage cannot be obtained or when spray drift may occur.

DO NOT apply aurally or through any type of irrigation system or equipment.

When spraying avoid overlapping, and shut off spray booms while starting, turning slowing and when stopped. Neglecting these procedures will cause higher rates of application resulting in crop damage. Delay spraying if heavy rain is expected within a few hours.

INSTRUCTIONS TO AVOID SPRAY DRIFT

DO NOT allow spray to drift onto adjacent turf sites or ornamental plants as even small amounts may injure sensitive plants. When drift may be a problem, take steps to reduce spray drift including:

DO NOT spray if wind is greater than 10 km/hr or if winds are gusting.

Use extreme caution when conditions are favourable for drift, i.e. high temperatures and low relative humidity, especially when sensitive plants are located nearby. All plants not listed as turf species, under Crop section on this label should be considered as sensitive plants.

If sensitive plants are downwind, extreme caution must be used under all conditions. Drift from applications of this herbicide is likely to result in damage to sensitive plants adjacent to the treatment site.

DO NOT apply unless there is a 30 metre downwind buffer distance between the treated areas and native vegetation.

DO NOT apply unless there is a 10 metre downwind buffer distance between the area being sprayed and natural water bodies.

COMPATIBILITY

Agricrop Prolan 500 Herbicide is compatible with paraquat, glyphosate, oxyfluorfen and simazine.

RESISTANT WEEDS WARNING

GROUP	D	HERBICIDE
-------	----------	-----------

Agricrop Prolan 500 Herbicide is a member of the dinitroaniline group of herbicides. Agricrop Prolan 500 Herbicide has the inhibition of tubulin formation mode of action. For weed resistance management, Agricrop Prolan 500 Herbicide is a Group D herbicide.

Some naturally occurring weed biotypes resistant to the product Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Agricrop Prolan 500 Herbicide or other Group D herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Agricrop Pty. Ltd accepts no liability for any losses that may result from the failure of Agricrop Prolan 500 Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture or Agricrop representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

When applying spray, avoid contact with leaves and other parts of trees and vines.

Do not apply under weather conditions or from spraying equipment, that may result in chemical movement by spray drift or storm run-off drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT contaminate dams, rivers, drains or streams with chemical or used container.

STORAGE AND DISPOSAL

Keep out of reach of children.

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver **empty** packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale spray mist.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

MSDS

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE

The use of Agricrop Prolan 500 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Agricrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Agricrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

In an Emergency Dial 000 Police or Fire Brigade

Batch Number:

Date of Manufacture: